



	Acetaminophen Examples: Tylenol, PediaCare, Little Fevers		Ibuprofen Examples: Motrin, Advil			Diphenhydramine Example: Benadryl	
<b>Frequency that can be given:</b>	Every 4-6 hrs		Every 6-8 hrs			Every 6-8 hrs	
<b>Name on bottle:</b>	** Infant's and Children's "Unconcentrated" Liquid/Suspension	Chewable Tablets	Infants "Concentrated"	Infant's and Children's Liquid/Suspension	"Children's" Chewable Tablets	Liquid/Suspension	Meltaway Tablets
<b>Concentration:</b>	160MG/5ML	1 Tablet = 80MG	50MG/1.25ML	100MG/5ML	1 Tablet = 50MG	12.5MG/5ML	1 Tablet = 12.5MG
<b>Age:</b>	2 Months or older		6 Months or older			6 Months or older	
<b>Weight:</b>	12-15 lbs	2.5 ML	-	1.25 ML	2.5ML	-	2.5 ML
	15-19 lbs	3 ML	-	1.5 ML	3 ML	-	2.5 ML
	19-24 lbs	4 ML	-	2 ML	4 ML	-	3.75 ML
	24-30 lbs	5 ML	2 Tablets	2.5 ML	5 ML	2 Tablets	5 ML
	30-40 lbs	6 ML	2 ½ Tablets	3.5 ML	7.5 ML	3 Tablets	6.25 ML
	38-48 lbs	7.5 ML	3 Tablets	4.25	8.5 ML	3 ½ Tablets	7.5 ML
	46-56 lbs	9 ML	3 ½ Tablets	-	10 ML	4 Tablets	10 ML
							2 Tablets

- \*\* = Tylenol labeling will say infants Suspension OR Children's Suspension = **THEY ARE THE SAME PRODUCT** because the concentration (160 MG = 5 ML) of the Tylenol is the same. This product will either come with a "syringe" or a "medicine cup". We always recommend using a syringe as it is more precise.
- **KEY POINTS ABOUT THESE PRODUCTS:** **ALWAYS** review the dosage on the bottle/box as the concentrations may vary among brand/manufacturer. The concentration/drug information can be found on the box where the "drug facts" are listed. A lot of the medications will say "talk to your doctor first before giving" – if we have recommended the medication and you use the correct dose according to this chart, it is OK to use. **We recommend bringing the medication that you will be using to your visit for verification. IF YOU MAKE A MISTAKE/ACCIDENTILY GIVE THE WRONG MEDICATION – CALL POISON CONTROL @ 1800-222-1222**